

AD-A215 217

FILE COPY


REPORT DOCUMENTATION PAGE			Form Approved OMB No. 0704-0188	
<small>Public reporting burden for this collection of information is estimated to average 1 hour per response, including the time for reviewing instructions, searching existing data sources, gathering and maintaining the data needed, and completing and reviewing the collection of information. Send comments regarding this burden estimate or any other aspect of this collection of information, including suggestions for reducing this burden, to Washington Headquarters Services, Directorate for Information Operations and Reports, 1215 Jefferson Davis Highway, Suite 1204, Arlington, VA 22202-4302, and to the Office of Management and Budget, Paperwork Reduction Project (0704-0188), Washington, DC 20503.</small>				
1. AGENCY USE ONLY (Leave blank)	2. REPORT DATE 1980	3. REPORT TYPE AND DATES COVERED Final		
4. TITLE AND SUBTITLE RATIONAL APPROXIMATIONS WITH HANKEL-NORM CRITERION		5. FUNDING NUMBERS  PE61102F 2304/A6		
6. AUTHOR(S) Y. Genin and S. Kung				
7. PERFORMING ORGANIZATION NAME(S) AND ADDRESS(ES) Stanford University Department of Electrical Engineering Stanford CA 94305		8. PERFORMING ORGANIZATION REPORT NUMBER  AFOSR-TR-89-1609		
9. SPONSORING/MONITORING AGENCY NAME(S) AND ADDRESS(ES) AFOSR BLDG 410 BAFB DC 20332-6448		10. SPONSORING/MONITORING AGENCY REPORT NUMBER  F49620-79-C-0058		
11. SUPPLEMENTARY NOTES				
12a. DISTRIBUTION/AVAILABILITY STATEMENT  Approved for public release; distribution unlimited.		12b. DISTRIBUTION CODE  in this report		
13. ABSTRACT (Maximum 200 words) A two-variable approach to the model reduction problem with Hankel-norm criterion is discussed. The problem is proved to be reducible to obtain a two-variable all-pass rational function, interpolating a set of parametric values at specified points inside the unit circle. A polynomial formulation and the properties of the optimal Hankel norm approximations are then shown to result directly from the general form of the solution of the interpolation problem considered. As a consequence, the recursive Nevanlinna algorithm can be employed and the essential stability properties of the solution can be established with the help of the Nevanlinna matrix (9). This short paper is meant to briefly summarize the work in the full paper (8), where the reader is referred to for more details.				
14. SUBJECT TERMS			15. NUMBER OF PAGES 2	
			16. PRICE CODE	
17. SECURITY CLASSIFICATION OF REPORT unclassified	18. SECURITY CLASSIFICATION OF THIS PAGE unclassified	19. SECURITY CLASSIFICATION OF ABSTRACT	20. LIMITATION OF ABSTRACT	

NSN 7540-01-280-5500

89 12 05 063

Standard Form 298 (890104 Draft)  
Prescribed by ANSI Std. Z39-18  
200-01

# DISCLAIMER NOTICE



THIS DOCUMENT IS BEST QUALITY AVAILABLE. THE COPY FURNISHED TO DTIC CONTAINED A SIGNIFICANT NUMBER OF PAGES WHICH DO NOT REPRODUCE LEGIBLY.

# RATIONAL APPROXIMATIONS WITH HANKEL-NORM CRITERION\*

Y. Genin\*\*  
Philips Research Lab.  
2, avenue van Becelaere  
B-1170 Brussels, Belgium

S. Kung  
Dept. of Electrical Engineering - Systems  
University of Southern California  
Los Angeles, CA 90007

## Abstract

A two-variable approach to the model reduction problem with Hankel norm criterion is discussed. The problem is proved to be reducible to obtain a two-variable all-pass rational function, interpolating a set of parametric values at specified points inside the unit circle. A polynomial formulation and the properties of the optimal Hankel norm approximations are then shown to result directly from the general form of the solution of the interpolation problem considered. As a consequence, the recursive Nevanlinna algorithm can be employed and the essential stability properties of the solution can be established with the help of the Nevanlinna matrix [9]. This short paper is meant to briefly summarize the work in the full paper [8], where the reader is referred to for more details.

## Introduction

In 1971, Adanjan, Arov and Krain [1] have obtained various powerful results related to the theory of bounded Hankel operators, which have great significance in the model reduction problem. Although their somewhat abstract original formulation in the framework of operator theory has been hindering their fast dissemination into the engineering community, the importance of this contribution has begun to be recognized. The relevance of [1] to the model reduction problem was first mentioned by Kung [2] in 1972, while some numerical aspects of the question were published in [3]. In [4],[5], connections between the minimal Hankel-norm approximations and the balanced state-space realizations introduced by Moore are put into light and shown to lead to an optimal approximation algorithm, requiring the solution of two Lyapunov equations and a singular value decomposition. In [6],[7], Kung took a pure one-variable polynomial approach and proved that the optimal approximation problem can be reduced to a simple generalized eigenvalue formulation when the transfer function is available.

## A. Two-Variable Polynomial Formulation

The problem of rational approximations in Hankel-norm can be defined as follows. Let a transfer function

$h(z)$  or equivalently its impulse response be given and let  $g(z)$  be some rational approximation of  $h(z)$ . Since the coefficients of the impulse response of a stable linear system are the Markov parameters of its transfer function, comparing two impulse responses amounts to comparing two Hankel matrices; the Hankel norm is precisely the spectral norm of the difference of these two matrices and it has a close relationship with other conventional norms [6]. The problem of the optimal approximations can then be stated in two different forms [6],[7],[8]:

- for a preassigned tolerance, how to obtain a minimal order approximation of  $h(z)$
- for a given order, how to obtain the minimal norm approximation of  $h(z)$

Clearly, the above two problems are intimately related.

It turns out that the problem of optimal approximations in Hankel norm can be globally approached and solved via a two-variable polynomial formalism [8].

## A. Two-Variable Polynomial Formulation

Let  $h(z) = b(z)/a(z)$  be the original transfer function of order  $m$ .

$$\hat{a}(z) = z^n \bar{a}(1/\bar{z})$$

$r(\lambda, z), \pi(\lambda, z)$  be two two-variable polynomials of degree  $n$  in  $z$ .

$$\hat{r}(\lambda, z) = z^n \bar{r}(\bar{\lambda}, 1/\bar{z})$$

$\epsilon$  can be any constant of unit modulus.

and let us consider the following equation:

$$\lambda b(z)r(\lambda, z) - \epsilon \hat{a}(z)\hat{r}(\lambda, z) = \pi(\lambda, z)a(z) \quad (1)$$

The above equation clearly defines a linear system of equations in the unknown coefficients of the polynomials  $r(\lambda, z)$  and  $\pi(\lambda, z)$ . It can be proved [8] that its minimal degree solution  $r(\lambda, z), \pi(\lambda, z)$  enjoys the following properties. For any fixed real value of  $\lambda$  and with  $[\pi(\lambda, z)/\lambda r(\lambda, z)]$  standing for the stable projection of  $\pi(\lambda, z)/\lambda r(\lambda, z)$  (i.e., after deleting its non-strictly stable part).

- $[\pi(\lambda, z)/\lambda r(\lambda, z)]$  is a minimal order approximation of  $h(z)$ , whose Hankel norm does not exceed  $|\lambda|^{-1}$ .
- If  $\lambda_{s+1}^{-1}$  is the  $(s+1)^{th}$  singular value of the Hankel matrix associated with  $h(z)$  and if one has the strict inequality  $\lambda_s^{-1} > \lambda_{s+1}^{-1}$ ,

\* This research has been supported by the U.S. Army Research Office, Contract DAAG-29-79-C-0215, the Air Force Office of Scientific Research, A.F. Systems Command, Contract AF 44-620-79-C-0058 and the National Science Foundation under Grant NSF-ENG-78-10003 and Grant NSF-ENG-73-28573. F49620-79-C-0058  
\*\* Currently on leave at the Information Systems Laboratory, Stanford University, CA 94303.

$\{r(\lambda_{s+1}, z)/r(\lambda_{s+1}, z)\}$  is the minimal Hankel norm approximation of order  $s$  of  $h(z)$  and the value of the norm is precisely  $\lambda_{s+1}$ .

## B. Parametric All-Pass Rational Interpolation

The above results can be easily proved with the help of standard matrix and polynomial techniques by a mere reformulation of the problem [8]. Let us indeed rewrite (1) in the following form:

$$\lambda \frac{b(z)}{a(z)} - \frac{r(\lambda, z)}{r(\lambda, z)} = \frac{r(\lambda, z)a(z)}{r(\lambda, z)a(z)} \quad (2)$$

Let us assume the zeros of  $a(z)$  to be distinct ( $z_1, z_2, \dots, z_n$ ) for the sake of brevity. With the notation  $u_i = b(z_i)/a(z_i)$ , solving (1) is clearly equivalent to obtain the minimal degree function  $s(\lambda, z) = r(\lambda, z)/r(\lambda, z)$  interpolating the parametric values  $u_i$  at the points  $z_i$ , i.e.,  $s(\lambda, z_i) = u_i$ . As a matter of fact, the latter problem can be solved [1], [8] by the so-called Nevanlinna algorithm [9], which solves the interpolation problem in a recursive way. As a result, the two-variable polynomial  $r(\lambda, z)$  can be constructed recursively and its stability properties investigated with the help of the Nevanlinna matrix, classically associated with the Nevanlinna algorithm [9]. Finally, the proof of the minimal order and minimal Hankel-norm properties of  $\{r(\lambda, z)/r(\lambda, z)\}$  is completed from a direct link existing between the Nevanlinna matrix and the singular values of the Hankel matrix associated with  $h(z)$ . [1], [8].

## References

- [1] V.M. Adamjan, D.Z. Arov, M.G. Krein, "Analytic properties of Schmidt pairs for a Hankel operator and the generalized Schur-Takagi problem," Math. USSR Sbornik, 15, pp. 31-73, 1971.
- [2] S. Kung, "A new identification and model reduction algorithm via singular value decompositions," Proc. of the 12th Asilomar Conference on Circuits, Systems and Computers, Nov. 6-8, 1978. Submitted to IEEE Trans. on AC.
- [3] A. Bultheel, "Numerieke aspecten van de Adamjan-Arov-Krein benadering," Second part of doctoral thesis, Faculty of Sciences, Leuven University, Belgium, 1979.
- [4] L.M. Silverman and Mamar Bettayeb, "Optimal approximation of linear systems," Proceedings, 1980 JACC, San Francisco, CA, Aug. 13-15, 1980.
- [5] A. Bultheel, P. Dewilde, "On the Adamjan-Arov-Krein approximation, identification and balanced realization," submitted to IEEE Trans. on CAS.
- [6] S. Kung, "Optimal Hankel-norm model reduction - scalar systems," Proceedings, 1980 JACC, San Francisco, CA, Aug. 13-15, 1980.
- [7] S. Kung and D.W. Lin, "Optimal Hankel-norm model reduction - matrix systems," Proceedings, IEEE 19th CDC, Albuquerque, NM, Dec. 10-12, 1980.
- [8] Y. Genin and S. Kung, "A two-variable approach to the model reduction problem with Hankel-norm criterion," submitted to IEEE Trans. on CAS.
- [9] R. Nevanlinna, "Über beschränkte analytische Funktionen," Ann. Acad. Sci. Fenn. A-32, pp. 1-75, 1929.

Accession For		
NTIS GRA&I	<input checked="" type="checkbox"/>	
DTIC TAB	<input type="checkbox"/>	
Unannounced	<input type="checkbox"/>	
Justification		
By		
Distribution/		
Availability Codes		
Dist	Avail and/or Special	
A-1	20	Circ-23